


Accepted Manuscript

Title: Therapeutic time window of anti-high mobility group box-1 antibody administration in mouse model of spinal cord injury

Authors: Masahide Nakajo, Naohiro Uezono, Hideyuki Nakashima, Hidenori Wake, Setsuro Komiya, Masahiro Nishibori, Kinichi Nakashima



PII: S0168-0102(18)30125-1
DOI: <https://doi.org/10.1016/j.neures.2018.03.004>
Reference: NSR 4160

To appear in: *Neuroscience Research*

Received date: 1-3-2018
Revised date: 19-3-2018
Accepted date: 19-3-2018

Please cite this article as: Nakajo, Masahide, Uezono, Naohiro, Nakashima, Hideyuki, Wake, Hidenori, Komiya, Setsuro, Nishibori, Masahiro, Nakashima, Kinichi, Therapeutic time window of anti-high mobility group box-1 antibody administration in mouse model of spinal cord injury. *Neuroscience Research* <https://doi.org/10.1016/j.neures.2018.03.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Therapeutic time window of anti-high mobility group box-1 antibody administration in mouse model of spinal cord injury

Author Affiliations:

Masahide Nakajo^{a,b}, Naohiro Uezono^{a,b}, Hideyuki Nakashima^a, Hidenori Wake^c, Setsuro Komiya^b, Masahiro Nishibori^c & Kinichi Nakashima^a

^aDepartment of Stem Cell Biology and Medicine, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maedashi, Higashi-ku, Fukuoka 812-8581, Japan.

^bDepartment of Orthopaedic Surgery, Graduate School of Medical and Dental Sciences, Kagoshima University, 8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan.

^cDepartment of Pharmacology, Graduate School of Medicine, Dentistry, and Pharmaceutical Sciences, Okayama University, 2-5-1 Shikata-cho, Kita-ku, Okayama 700-8558, Japan.

Highlights

- Administration of anti-HMGB1 mAb within 6 h after SCI improves functional recovery
- Anti-HMGB1 mAb protects the BSCB
- Anti-HMGB1 mAb suppresses inflammation after SCI

ABSTRACT

Spinal cord injury (SCI) is a devastating neurologic disorder that often leads to

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات