

Accepted Manuscript

Phase-locking and bistability in neuronal networks with synaptic depression

Zeynep Akcay, Xinxian Huang, Farzan Nadim, Amitabha Bose

PII: S0167-2789(17)30218-X
DOI: <https://doi.org/10.1016/j.physd.2017.09.007>
Reference: PHYSD 31949

To appear in: *Physica D*

Received date: 17 April 2017
Revised date: 22 August 2017
Accepted date: 26 September 2017

Please cite this article as: Z. Akcay, X. Huang, F. Nadim, A. Bose, Phase-locking and bistability in neuronal networks with synaptic depression, *Physica D* (2017), <https://doi.org/10.1016/j.physd.2017.09.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Phase-locking and bistability in neuronal networks with synaptic depression

Zeynep Akcay¹, Xinxian Huang², Farzan Nadim^{2,3}, Amitabha Bose²

1. Department of Mathematics and Computer Science, Queensborough Community College, Bayside, NY 11364, USA
2. Department of Mathematical Sciences, New Jersey Institute of Technology, Newark, NJ, 07102, USA
3. Federated Department of Biological Sciences, New Jersey Institute of Technology and Rutgers University, Newark, NJ 07102, USA

Corresponding Author: Amitabha Bose; bose@njit.edu

Abstract

We consider a recurrent network of two oscillatory neurons that are coupled with inhibitory synapses. We use the phase response curves of the neurons and the properties of short-term synaptic depression to define Poincaré maps for the activity of the network. The fixed points of these maps correspond to phase-locked modes of the network. Using these maps, we analyze the conditions that allow short-term synaptic depression to lead to the existence of bistable phase-locked, periodic solutions. We show that bistability arises when either the phase response curve of the neuron or the short-term depression profile changes steeply enough. The results apply to any Type I oscillator and we illustrate our findings using the Quadratic Integrate-and-Fire and Morris-Lecar neuron models.

Keywords: Coupled Oscillators, Phase Response Curve, Two-dimensional Poincaré Map, Bistability, Short-Term Synaptic Depression

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات