

Title: Prenatal exposure to valproic acid alters the development of excitability in the postnatal rat hippocampus

Authors: Yukiko Fueta, Yuko Sekino, Sachiko Yoshida, Yasunari Kanda, Susumu Ueno

PII: S0161-813X(18)30003-2
DOI: https://doi.org/10.1016/j.neuro.2018.01.001
Reference: NEUTOX 2279

To appear in: *NEUTOX*

Received date: 5-12-2016
Revised date: 3-12-2017
Accepted date: 4-1-2018

Please cite this article as: Fueta Yukiko, Sekino Yuko, Yoshida Sachiko, Kanda Yasunari, Ueno Susumu.Prenatal exposure to valproic acid alters the development of excitability in the postnatal rat hippocampus.*Neurotoxicology* https://doi.org/10.1016/j.neuro.2018.01.001

Prenatal exposure to valproic acid alters the development of excitability in the postnatal rat hippocampus

Yukiko FUETA[a], Yuko SEKINO[b,c], Sachiko YOSHIDA[d], Yasunari KANDA[b], Susumu UENO[e,*]

[a]Department of Environmental Management and Control, School of Health Sciences, University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan.

[b]Division of Pharmacology, National Institute of Health Sciences, Kamiyoga 1-18-1, Setagaya-ku, Tokyo 158-8501, Japan.

[c]Laboratory of Chemical Pharmacology, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 7-3-1 Hongo Bunkyo-ku, Tokyo 113-0033, Japan.

[d]Department of Environment and Life Sciences, Toyohashi University of Technology, 1-1 Hibarigaoka, Tempaku-cho, Toyohashi, Aichi 441-8580, Japan

[e]Department of Occupational Toxicology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu 807-8555, Japan.