

Available online at www.sciencedirect.com

ScienceDirect

Ann. I. H. Poincaré – AN ●●● (●●●●) ●●●●●●

ANNALES
DE L'INSTITUT
HENRI
POINCARÉ
ANALYSE
NON LINÉAIREwww.elsevier.com/locate/anihpc

On the asymptotic growth of positive solutions to a nonlocal elliptic blow-up system involving strong competition [☆]

Susanna Terracini ^{*}, Stefano Vita*Dipartimento di Matematica "Giuseppe Peano", Università di Torino, Via Carlo Alberto, 10, 10123 Torino, Italy*

Received 23 June 2016; received in revised form 10 July 2017; accepted 30 August 2017

Abstract

For a competition-diffusion system involving the fractional Laplacian of the form

$$-(-\Delta)^s u = uv^2, \quad -(-\Delta)^s v = vu^2, \quad u, v > 0 \text{ in } \mathbb{R}^N,$$

with $s \in (0, 1)$, we prove that the maximal asymptotic growth rate for its entire solutions is $2s$. Moreover, since we are able to construct symmetric solutions to the problem, when $N = 2$ with prescribed growth arbitrarily close to the critical one, we can conclude that the asymptotic bound found is optimal. Finally, we prove existence of genuinely higher dimensional solutions, when $N \geq 3$. Such problems arise, for example, as blow-ups of fractional reaction-diffusion systems when the interspecific competition rate tends to infinity.

© 2017 Published by Elsevier Masson SAS.

MSC: primary 35J65; secondary 35B40, 35B44, 35R11, 81Q05, 82B10

Keywords: Fractional Laplacian; Spatial segregation; Strongly competing systems; Entire solutions

1. Introduction and main results

This paper deals with the existence and classification of positive entire solutions to polynomial systems involving the (possibly) s -fractional Laplacian of the following form:

$$-(-\Delta)^s u = uv^2, \quad -(-\Delta)^s v = vu^2, \quad u, v > 0 \text{ in } \mathbb{R}^N.$$

Such systems arise, for example, as blow-ups of fractional reaction-diffusion systems when the interspecific competition rate tends to infinity. In this framework, the existence and classification of entire solutions plays a key role in the

[☆] Work partially supported by the ERC Advanced Grant 2013 n. 339958 *Complex Patterns for Strongly Interacting Dynamical Systems – COMPAT* and by the PRIN-2012-74FYK7 Grant *Variational and perturbative aspects of nonlinear differential problems*. The authors wish to thank Alessandro Zilio for many fruitful conversations.

^{*} Corresponding author.

E-mail addresses: susanna.terracini@unito.it (S. Terracini), stefano.vita@unito.it (S. Vita).

asymptotic analysis (see, for instance, [15,17]). The case of standard diffusion ($s = 1$) has been intensively treated in the recent literature, also in connection with a De Giorgi-like conjecture about monotone solutions being one dimensional. In particular, a complete classification of solutions having linear growth (the lowest possible growth rate) has been given in [1,2,7–9,16,20]. On the other hand, when $s = 1$, positive solutions having arbitrarily large polynomial growth were discovered in [2] and with exponential growth in [14].

Competition-diffusion nonlinear systems with k -components involving the fractional Laplacian have been the object of a recent literature, starting with [18,19], where the authors provided asymptotic estimates for solutions to systems of the form

$$\begin{cases} (-\Delta)^s u_i = f_{i,\beta}(u_i) - \beta u_i \sum_{j \neq i} a_{ij} u_j^2, & i = 1, \dots, k, \\ u_i \in H^s(\mathbb{R}^N), \end{cases} \tag{1.1}$$

where $N \geq 2$, $a_{ij} = a_{ji} > 0$, when $\beta > 0$ (the competition parameter) goes to $+\infty$. Moreover we consider $f_{i,\beta}$ as continuous functions which are uniformly bounded on bounded sets with respect to β (see [18,19] for details). The fractional Laplacian is defined for every $s \in (0, 1)$ as

$$(-\Delta)^s u(x) = c(N, s) \text{PV} \int_{\mathbb{R}^N} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy.$$

In order to state our results, we adopt the approach of Caffarelli–Silvestre [5], and we see the fractional Laplacian as a Dirichlet-to-Neumann operator; that is, we consider the extension problem for (1.1). In other words, we study an auxiliary problem in the upper half space in one more dimension¹; that is, letting $a := 1 - 2s$, for any $i = 1, \dots, k$ the localized version of (1.1),

$$\begin{cases} L_a u_i = 0, & \text{in } B_1^+ \subset \mathbb{R}_+^{N+1}, \\ \partial_y^a u_i = f_{i,\beta}(u_i) - \beta u_i \sum_{j \neq i} a_{ij} u_j^2, & \text{in } \partial^0 B_1^+ \subset \partial \mathbb{R}_+^{N+1} = \mathbb{R}^N \times \{0\}, \end{cases} \tag{1.2}$$

where the degenerate/singular elliptic operator L_a is defined as

$$-L_a u := \text{div}(y^a \nabla u),$$

and the linear operator ∂_y^a is defined as

$$-\partial_y^a u := \lim_{y \rightarrow 0^+} y^a \frac{\partial u}{\partial y}.$$

The new problem (1.2) is equivalent to the original one when we deal with solutions in the energy space associated with the two operators. In fact a solution U to the extension problem is the extension of the correspondent solution u of the original nonlocal problem in the sense that $U(x, 0) = u(x)$. Let us remark that if $s = \frac{1}{2}$, then $a = 0$ and hence $L_0 = -\Delta$ and the boundary operator $-\partial_y^0$ becomes the usual normal derivative ∂_y . Moreover we remark that the extension problem has a variational nature in some weighted Sobolev spaces related to the Muckenhoupt A_2 -weights (see for instance [10]). Hence, given $\Omega \subset \mathbb{R}_+^{N+1}$, we can introduce the Hilbert spaces

$$H^{1;a}(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} : \int_{\Omega} y^a (|u|^2 + |\nabla u|^2) < +\infty \right\},$$

and

$$H_{loc}^{1;a}(\overline{\mathbb{R}_+^{N+1}}) := \left\{ u : \mathbb{R}_+^{N+1} \rightarrow \mathbb{R} : \forall r > 0, u|_{B_r^+} \in H^{1;a}(B_r^+) \right\},$$

¹ Throughout this paper we assume the following notations: $z = (x, y)$ denotes a point in \mathbb{R}_+^{N+1} , with $x \in \partial \mathbb{R}_+^{N+1} := \mathbb{R}^N$ and $y \in \mathbb{R}_+$. Moreover, $B_r^+(z_0) := B_r(z_0) \cap \mathbb{R}_+^{N+1}$ is the half ball, and its boundary is divided in the hemisphere $\partial^+ B_r^+(z_0) := \partial B_r^+(z_0) \cap \mathbb{R}_+^{N+1}$ and in the flat part $\partial^0 B_r^+(z_0) := \partial B_r^+(z_0) \setminus \partial^+ B_r^+(z_0)$. When the center of balls and spheres is omitted, then $z_0 = 0$.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات