



# Differentiating symptoms of social anxiety and depression in adults with social anxiety disorder

Brandon E. Gibb<sup>a,\*</sup>, Meredith E. Coles<sup>a</sup>, Richard G. Heimberg<sup>b</sup>

<sup>a</sup>*Department of Psychology, Binghamton University (SUNY), Binghamton, NY, 13902-6000, USA*

<sup>b</sup>*Adult Anxiety Clinic of Temple, Department of Psychology, Temple University, 1701 North 13th Street, Philadelphia, PA 19122-6085, USA*

Received 17 September 2003; received in revised form 20 July 2004; accepted 5 August 2004

---

## Abstract

Although studies have suggested a strong overlap between social anxiety disorder and depression, this is the first study to examine the ability of commonly used measures to differentiate symptoms of these disorders in a sample of clients with social anxiety disorder. Structural equation modeling revealed that commonly used measures of social anxiety and depression can differentiate the two constructs, rather than simply reflecting a single construct of overall distress. Logistic regression analyses indicated that scores on depressive symptom measures could predict which socially anxious clients met criteria for a comorbid depressive disorder.

© 2004 Elsevier Ltd. All rights reserved.

*Keywords:* Anxiety; Depression; Comorbidity; Structural equation modeling

---

## 1. Introduction

A number of studies have noted the lack of discriminant validity traditionally found among measures of anxiety and depression (for a review, see [Gotlib & Cane, 1989](#)). Given this, [Clark and Watson \(1991\)](#) proposed the tripartite model of anxiety

---

\*Corresponding author. Tel.: +1-607-770-2511; fax: +1-607-777-3665.

E-mail address: [bgibb@binghamton.edu](mailto:bgibb@binghamton.edu) (B.E. Gibb).

and depression in which they predicted that certain symptom clusters that would be relatively unique to anxiety and to depression as well as symptoms that would be common to both disorders or emotional states. Specifically, they suggested that both anxiety and depression would be characterized by negative affect and that symptoms of physiological arousal would be relatively unique to anxiety whereas symptoms of anhedonia or low positive affect would be relatively unique to depression. Despite general support for the tripartite model (e.g., Brown, Chorpita, & Barlow, 1998; Joiner, 1996; Joiner, Cantanzaro, & Laurent, 1996; Watson et al., 1995a, b), there is evidence that social anxiety disorder, as well as depression, may be characterized by anhedonia (Brown et al., 1998; Watson, Clark, & Carey, 1988), suggesting that measures of anhedonia or low positive affect may not differentiate the two disorders.

Given the high degree of symptom and diagnostic comorbidity between social anxiety and depression (Alpert et al., 1999; Alpert, Maddocks, Rosenbaum, & Fava, 1994; Kessler et al., 1996; Kessler, Stang, Wittchen, Stein, & Walters, 1999; Schneier, Johnson, Hornig, Liebowitz, & Weissman, 1992; Van Ameringen, Mancini, Styan, & Donison, 1991; Zimmerman, McDermut, & Mattia, 2000), it is important for researchers to be able to validly discriminate between the two disorders. Specifically, it is important to determine whether measures of social anxiety and depressive symptoms assess the constructs of interest rather than simply assessing shared symptoms of negative affect.

The goal of this study, therefore, was to determine whether commonly used measures of social anxiety and depression can be used to differentiate the two constructs in a sample of clients with social anxiety disorder. Specifically, we used structural equation modeling to test a model in which social anxiety and depression were specified as separate constructs versus one in which they were specified as a single construct, representing overall general distress. We then examined the ability of the depressive symptom measures to predict which clients met criteria for a comorbid depressive disorder.

## 2. Method

### 2.1. Participants

The sample consisted of 113 clients who sought treatment for performance and/or interpersonal anxiety. All clients were between the ages of 18 and 65 ( $M = 36.5$ ,  $SD = 9.8$ ) and 58 (51.3%) were female. Clients were recruited through advertisements in local newspapers and flyers. After completing a brief initial phone-screening interview, prospective clients were scheduled for a diagnostic interview. All participants in this study met *DSM-III-R* (American Psychiatric Association, 1987) criteria for a principal diagnosis of social anxiety disorder after a semi-structured interview with the Anxiety Disorders Interview Schedule-Revised (ADIS-R; DiNardo & Barlow, 1988). ADIS-R interviews were conducted by advanced graduate students and post-doctoral fellows trained to reliability standards (DiNardo, Moras, Barlow, Rapee, & Brown, 1993). Clinicians using the ADIS-R

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات