



The influence of age on the distribution of self-rated health, burnout and their associations with psychosocial work conditions

Yawen Cheng ^{a,*}, I-Shin Chen ^a, Chiou-Jong Chen ^b, Hermann Burr ^c, Hans Martin Hasselhorn ^c

^a Institute of Health Policy and Management, College of Public Health, National Taiwan University, Taiwan

^b Institute of Occupational Safety and Health, Council of Labor Affairs, Taipei, Taiwan

^c Federal Institute for Occupational Safety and Health, Germany

ARTICLE INFO

Article history:

Received 30 October 2012

Received in revised form 11 December 2012

Accepted 13 December 2012

Keywords:

Age

Self-rated health

Burnout

Psychosocial work conditions

ABSTRACT

Objective: Self-rated health (SRH) and burnout are commonly used health indicators. This study was designed to examine the age-specific patterns of SRH and burnout and their correlations with self-reported disease symptoms, and to investigate the moderating effects of age on the associations of psychosocial work conditions with these two health measures.

Methods: Study subjects were 20,454 male and 16,875 female employees from 2 representative surveys conducted in 2007 and 2010 in Taiwan. SRH was assessed by a single item and burnout was assessed by the general burnout subscale of the Copenhagen Burnout Inventory (CBI). Psychosocial work conditions including working hours, psychological and physical job demands, job control, job insecurity and workplace justice were assessed by a validated questionnaire.

Results: Older workers were at higher risk for poor SRH which was correlated with the presence of multiple disease symptoms. In contrast, employees aged 30–40 years had higher burnout which was strongly correlated with emotional disturbance. Among all the examined work factors, low workplace justice was found to be the most predominant predictor for poor SRH and high burnout, and the associations were more profound in younger workers than in older workers.

Conclusion: SRH and burnout were differentially related to age, and there were evidences of moderating effects of age on the associations between psychosocial work conditions and health. Researchers and health practitioners should pay attention to the influence of age when using the measures of SRH and burnout as indicators to detect health risk in association with adverse psychosocial work conditions.

© 2013 Elsevier Inc. All rights reserved.

Introduction

In epidemiologic studies concerning the health risks of psychosocial work hazards, self-rated health (SRH) and burnout are commonly used outcome indicators. The single-item measure for SRH is known to be a composite indicator for all dimensions of health. On the other hand, burnout is conceptualized as an affective reaction to prolonged stress, and is reflected primarily in emotional exhaustion, physical fatigue and cognitive weariness [1,2].

The measures of SRH and burnout have been found to have high predictive values, and have been suggested to be useful tools in identifying employees who are at high risk for sickness absence, cardiovascular disease, musculoskeletal disorders and other physical and mental diseases [3–15]. In occupational settings, there have been an enormous amount of studies linking negative psychosocial work characteristics, including high job demands, lower job control, poor

social support at work, precarious employment, effort-reward imbalance and organizational injustice, to poor SRH and high level of burnout [16–22].

However, to our knowledge, few studies have explicitly investigated the potential moderating effects of age on the associations between psychosocial work conditions and health risks [23]. It is possible that similarly stressful work conditions such as long working hours and heavy workloads may induce greater physiological responses in older workers, due to an increased susceptibility associated with ageing. On the other hand, lack of job control or employment insecurity might cause greater psychological distress among younger workers, taking in account that work is supposed to play a central role in the earlier stage of working life.

Moreover, while previous studies have suggested that older people, who in general have greater degree of health problems, may evaluate their health differently from younger people who are mostly free of serious disease [24,25], the influence of age on the distribution of SRH and burnout has rarely been studied in general employees across the full age range. It is reasonable to assume that a person's general assessment of his or her health is reciprocally influenced by the status of mental and physical exhaustion. For instance, a recent study of a

* Corresponding author at: Institute of Health Policy and Management, College of Public Health, National Taiwan University, 17, Xu-Zhou Rd., Taipei, Taiwan. Tel.: +886 2 3366 8056; fax: +886 2 2321 5048.

E-mail address: ycheng@ntu.edu.tw (Y. Cheng).

sample of air force personnel indicated that perceived health was highly correlated with burnout—the latter was defined by the level of depletion of one's intrinsic energetic resources [26]. Nevertheless, information with regard to the correlations between self-rated health and burnout, as well as their correlations with the presence of disease symptoms—either physical or psychological—has been limited.

As illustrated in Fig. 1, this study was designed to examine the influence of age on the distribution of SRH and burnout, and to explore the moderating effects of age on the associations of psychosocial work conditions with SRH and burnout. Data from 2 rounds of national surveys encompassing employees of the full age range were utilized for this study.

Various instruments have been developed to measure burnout. Among all, the Maslach Burnout Inventory (MBI) has been the most widely used instrument. The MBI defines 'burnout' as a psychological consequence of prolonged stress arising out of work. According to the MBI, burnout is measured by a weighted sum of three components, namely 'emotional exhaustion', 'depersonalization' and 'reduced personal accomplishment' [1,27]. Similarly, the Oldenburg Burnout Inventory (OLBI) defines and assesses 'burnout' by the levels of 'exhaustion' and 'disengagement' [15,28]. By contrast, some measures, for instance, the Schirom Melamed Burnout Questionnaire and the personal burnout scale of the Copenhagen Burnout Inventory (CBI) [29,30], focus on the degree of general physical and psychological exhaustion experienced by the person regardless of its relation to work.

In this study, we focused on the core component of burnout, i.e., the overall physical and psychological exhaustion. We deliberately chose the personal burnout scale of the CBI which does not attribute burnout to work or other specific causes, because the association between work conditions and burnout was one of the study objectives to be examined.

Methods

Study subjects and study design

The Council of Labour Affairs of Taiwan has conducted nation-wide cross-sectional surveys in an interval of 3 to 5 years since 1988. The surveys are designed to assess multiple dimensions of safety and health issues in the work population. In these surveys, subjects were selected through a two-stage random sampling process. In the first stage, all districts and villages throughout Taiwan were grouped into strata according to the level of urbanization. A random sample of districts and villages was chosen from each stratum. In the second stage, a random sample of households was selected within each district or village, and residents of the sampled households who were currently working at the time of survey were identified. Self-administered questionnaires

were delivered to the selected households by trained interviewers. Subjects were informed of the purpose of the survey and invited to participate in a face-to-face manner. After one week, questionnaires were collected and on-site checking was performed by the same interviewer to ensure the completeness.

Data for this study was drawn from the national surveys conducted in September 2007 and September 2010. In 2007, 22,476 employees were sampled. Among them, 19,329 subjects completed and returned questionnaires, with an overall response rate of 86%. In 2010, a total of 27,508 subjects (including employees, employers and self-employed workers) were sampled and 23,932 subjects completed and returned questionnaires, with an overall response rate of 87%. For this study, we restricted analyses to employees aged between 20 and 65 years old. This resulted in a pooled sample of altogether 37,329 employees (20,454 men and 16,875 women).

Measures of health status and burnout

Self-rated health was assessed by a single-item question "In general, how is your health?", which has five possible answers ranging from "very good", "good", "moderate", "poor" and "very poor". In this study, the responses were dichotomized into poor SRH (moderate, poor or very poor) vs. good SRH (very good or good).

Burnout status was assessed by the personal burnout subscale of the Chinese version of Copenhagen Burnout Inventory (CBI). With the CBI, burnout is conceptualized as a state of physical and psychological fatigue and exhaustion. Information regarding the English version and the Chinese version of the CBI, including the details of all subscales, items and calculation formula, can be found elsewhere [29,31]. The original scale contained 6 items: "How often do you feel tired?", "How often are you physically exhausted?", "How often are you emotionally exhausted?", "How often do you think: "I can't take it anymore?", "How often do you feel weak and susceptible to illness?" and "How often do you feel worn out?". The responses are coded in a 5-point scale ranging from "always"(score 100), "often" (75), "at times"(50), "not often"(25) and "never"(0). In the national surveys, however, one item—"How often do you feel worn out?"—was deleted due to the length constraint of our questionnaire. This item was chosen to be removed because its statement in Chinese appeared to be very similar to the item of "How often do you feel tired?". In the present study, the 5-item scale for personal burnout showed a high internal consistency, with Cronbach's alpha values of 0.9 for both men and women. A mean score of 50 or above was classified as high burnout in this study.

Participants were also asked if over the past 12 months they had experienced any of the following disease symptoms (yes/no): "musculoskeletal discomfort", "gastrointestinal irritation or illness",




Fig. 1. Influence of age on self-rated health, burnout and their associations with psychosocial work conditions.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات