



# The role of lateral occipital face and object areas in the face inversion effect

David Pitcher<sup>a,b,\*</sup>, Bradley Duchaine<sup>c</sup>, Vincent Walsh<sup>b</sup>, Galit Yovel<sup>d</sup>, Nancy Kanwisher<sup>a</sup>

<sup>a</sup> McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>b</sup> Institute of Cognitive Neuroscience, University College London, Alexandra House, 17 Queen Square, London WC1N 3AR, United Kingdom

<sup>c</sup> Department of Psychology and Brain Sciences, Dartmouth College, Hanover, NH 03755, USA

<sup>d</sup> Department of Psychology, Tel Aviv University, Tel Aviv 69978, Israel

## ARTICLE INFO

### Article history:

Received 31 January 2011

Received in revised form 19 August 2011

Accepted 22 August 2011

Available online 30 August 2011

### Keywords:

Face perception

Object recognition

Transcranial magnetic stimulation

Occipital face area

Lateral occipital area

## ABSTRACT

Stimulus inversion impairs face discrimination to a greater extent than discrimination of other non-face object categories. This finding has led to suggestions that upright faces are represented by mechanisms specialized for upright faces whereas inverted face representation depends on more general object recognition mechanisms. In the present study we tested the causal role of face-selective and object-selective cortical areas for upright and inverted face discrimination by transiently disrupting neural processing using transcranial magnetic stimulation (TMS). Participants matched upright and inverted faces while TMS was delivered over each participant's functionally localized right occipital face area (rOFA) or right lateral occipital area (rLO). TMS delivered over rOFA disrupted the discrimination of upright and inverted faces while TMS delivered over rLO impaired inverted face discrimination only. These results provide causal evidence that upright faces are represented by face-specific mechanisms whereas inverted faces are represented by both face-specific and object-specific mechanisms. The similar sensitivity of the OFA to upright and inverted faces is consistent with the hypothesis that the OFA processes facial features at an early stage of face processing.

© 2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

The face inversion effect (FIE) is a classic signature of face perception in which stimulus inversion disrupts face recognition more strongly than object recognition (Farah, Tanaka, & Drain, 1995; McKone, Kanwisher, & Duchaine, 2007; Yin, 1969). The FIE is taken as evidence for the existence of face-specific mechanisms in the brain that are tailored for processing upright faces only (Farah, 2004; Yin, 1969). However this account does not specify which mechanisms contribute to the perception of inverted faces and whether mechanisms that are not face-selective contribute to upright face recognition. Here we tested these questions using transcranial magnetic stimulation (TMS).

### 1.1. The neural basis of the face inversion effect

Neuropsychological studies of patients with impairments in visual processing provide causal evidence that upright and inverted faces are represented using distinct cognitive mechanisms (Farah, 2004; Yin, 1970). Some prosopagnosic patients are impaired (relative to control subjects) in recognizing upright but not inverted

faces (Farah, Wilson, Drain, & Tanaka, 1995). The same is true of developmental prosopagnosics, who have lifelong impairments in face recognition despite the absence of any known brain damage (Duchaine, Yovel, & Nakayama, 2007). By contrast, one patient with an object recognition impairment was normal at upright face recognition, but severely impaired in recognizing inverted faces, thus showing a face inversion effect that was many-fold larger in magnitude than that found in normal subjects (Moscovitch & Moscovitch, 2000; Moscovitch, Winocur, & Behrmann, 1997). These findings have been taken to show that inverted faces are processed through the generic object recognition pathway, whereas upright faces are processed in systems specialized for upright faces only.

Functional magnetic resonance imaging (fMRI) studies of neurologically normal participants have also examined how upright and inverted faces are represented in the brain. Several studies have demonstrated that the face-selective fusiform face area (FFA) (Kanwisher, McDermott, & Chun, 1997) exhibits a greater response to images of upright faces than to images of inverted faces (Epstein, Higgins, Parker, Aguirre, & Cooperman, 2006; Kanwisher, Tong, & Nakayama, 1998; Mazard, Schiltz, & Rossion, 2006; Yovel & Kanwisher, 2005; but see Aguirre, Singh, & D'Esposito, 1999; Haxby et al., 1999). By contrast, the scene-selective parahippocampal place area (PPA) (Epstein & Kanwisher, 1998) as well as the object recognition area in the lateral occipital cortex (LO) (Malach et al., 1995) exhibit greater responses to inverted faces than to upright faces (Aguirre et al., 1999; Epstein 2005; Haxby et al., 1999;

\* Corresponding author at: Laboratory of Brain and Cognition, National Institute for Mental Health, Bethesda, MD 20892, USA. Tel.: +1 301 435 4931.

E-mail addresses: [pitcherdj@mail.nih.gov](mailto:pitcherdj@mail.nih.gov), [dpitcher@mit.edu](mailto:dpitcher@mit.edu) (D. Pitcher).

Yovel & Kanwisher, 2005). These studies further demonstrate that upright and inverted faces are preferentially processed in functionally segregated and spatially distinct cortical areas.

In the present study we investigated the contribution of face-selective and object-selective cortical areas to upright and inverted face discrimination by combining the spatial precision of fMRI with the causal inferences one can draw from the neural disruption induced by transcranial magnetic stimulation (TMS). Participants performed a delayed match-to-sample discrimination task with upright and inverted faces while TMS was delivered over the functionally localized right occipital face area (rOFA) and right LO (rLO). The OFA is believed to play a role in early feature-based stages of face perception (Haxby et al., 1999) and TMS delivered over rOFA has been shown to disrupt discrimination of upright faces but not discrimination of non-face stimuli such as houses, objects, and human bodies (Pitcher, Charles, Devlin, Walsh, & Duchaine, 2009; Pitcher, Garrido, Walsh, & Duchaine, 2008; Pitcher, Walsh, Yovel, & Duchaine, 2007). The OFA has also been shown to exhibit a similar neural response to upright and inverted faces (Yovel & Kanwisher, 2005; but see Mazard et al., 2006) leading us to hypothesize that TMS delivered over the rOFA would disrupt both upright and inverted face discrimination. TMS delivered over rLO disrupts object discrimination (Chouinard, Whitwell, & Goodale, 2009) but not face discrimination (Pitcher et al., 2007, 2009). Based on the neuropsychological, fMRI and previous TMS evidence we predicted that TMS delivered over rLO would disrupt inverted face discrimination but have no effect on upright face discrimination.

## 2. Materials and methods

### 2.1. Participants

Ten right-handed participants with normal or corrected-to-normal vision (5 females, aged 19–27) gave informed consent as directed by the Massachusetts Institute of Technology IRB committee. No participants withdrew due to discomfort with TMS stimulation.

### 2.2. Materials

Closely matched face stimuli in which the component parts were altered were used and similar example stimuli are presented in Fig. 1 (Yovel & Kanwisher, 2004). These stimuli were used in a previous TMS study of the rOFA (Pitcher et al., 2007).

### 2.3. Procedure

The experiment used a two-by-three design in which participants discriminated upright and inverted faces while rTMS was delivered over rOFA, rLO or no TMS was delivered (the no TMS condition was included as a behavioural baseline). Fig. 2 shows the trial procedure. Stimuli were presented centrally on an SVGA 20 inch monitor (Resolution 1024 by 768, refresh rate 70 Hz). Participants sat 57 cm from the monitor with their heads stabilized in a chin rest and indicated by a right hand key press whether the sample stimulus was the same as the probe stimulus. They were instructed to respond accurately and as quickly as possible.

Face orientation (upright or inverted) was blocked and the order was balanced across participants (half of the participants started with upright faces, the other half with inverted faces). Three blocks of 40 trials (20 same trials, 20 different trials) were presented for each face orientation. During each block, rTMS was delivered over rOFA or rLO or no TMS was delivered. The order of TMS blocks was varied across participants and balanced using a Latin square design. Within each block the trial order was randomized. During the same testing session participants also completed a second TMS discrimination task using different stimuli that tested a different hypothesis.

### 2.4. Imaging

TMS target sites (rOFA and rLO) were individually identified in all participants using a standard fMRI localizer task (Pitcher, Dilks, Saxe, Triantafyllou, & Kanwisher, 2011). Functional data were acquired over 4 blocked-design functional runs lasting 234 s each. Scanning was performed in a 3.0 T Siemens Trio scanner at the A. A. Martinos Imaging Center at the McGovern Institute for Brain Research at the Massachusetts Institute of Technology. Functional images were acquired with a Siemens 32-channel phased array head-coil and a gradient-echo EPI sequence (32 slices, repetition time (TR) = 2 s, echo time = 30 ms, voxel size = 3 mm × 3 mm × 3 mm, and 0.6 mm interslice gap) providing whole brain coverage (slices were aligned with

the anterior/posterior commissure). In addition, a high-resolution T-1 weighted MPRAGE anatomical scan was acquired for anatomically localizing the functional activations. Each functional run contained two sets of five consecutive dynamic stimulus blocks (faces, bodies, scenes, objects or scrambled objects) sandwiched between rest blocks, to make two blocks per stimulus category per run. Each block lasted 18 s and contained stimuli from one of the five stimulus categories.

Data were analyzed with FS-FAST, Freesurfer (<http://surfer.nmr.mgh.harvard.edu/>) (Dale, Fischl, & Sereno, 1999; Fischl, Sereno, & Dale, 1999). Before statistical analysis, images were motion corrected (Cox & Jesmanowicz, 1999), smoothed (3 mm FWHM Gaussian kernel), detrended, and fit using a gamma function ( $\Delta = 2.25$  and  $\tau = 1.25$ ). The pre-processing did not involve any spatial normalization of subjects in a common reference space (e.g., Talairach transformations). The functional data of each subject were co-registered with that subject's anatomical image.

Significance maps of the brain were computed using the same statistical threshold for both TMS target sites ( $p = 10^{-4}$ , uncorrected) (see Fig. 3). The rOFA was identified using a contrast of dynamic faces greater than dynamic objects and was always located on the lateral surface of the occipital lobe posterior to the face-selective rFFA (mean MNI co-ordinates = 43, -79, -11). The rLO was identified using a contrast of dynamic objects greater than scrambled objects and was always located on the lateral surface of the occipital lobe and was superior to the rOFA (mean MNI co-ordinates = 44, -73, -6). The coordinates and strength of the peak responses varied across participants but rOFA and rLO were identified in each participant.

### 2.5. TMS stimulation and site localization

TMS target sites were individually identified using the Brainsight TMS-MRI co-registration system (Rogue Research, Montreal, Canada), utilizing individual high resolution structural and functional MRI scans for each participant. The rOFA and rLO were localized by overlaying individual activation maps from the fMRI localizer task for the face and object analysis and identifying the voxel exhibiting the greatest activation in each category-selective region. The surface coil locations were then marked on each participant's head. To ensure accurate coil placement during the experiment the position of the coil was tracked and monitored during half of the TMS blocks using the Brainsight system.

TMS was delivered at 10 Hz and 60% of maximal stimulator output, using a Magstim Super Rapid Stimulator (Magstim, UK) and a 70 mm figure-of-eight coil, with the coil handle pointing upwards and parallel to the midline. A single intensity was used on the basis of previous studies (O'Shea, Muggleton, Cowey, & Walsh, 2004; Silvano, Lavie, & Walsh, 2005) and for ease of comparison with similar studies of the rOFA and rLO (Pitcher et al., 2007, 2009). In TMS blocks, TMS was delivered at a frequency of 10 Hz for 500 ms and its onset coincided with the onset of the test stimulus.

## 3. Results

Accuracy was measured with  $d'$  (Green & Swets, 1966), an unbiased measure of discrimination performance, and the mean data are shown in Fig. 4a. A two-by-three repeated measures analysis of variance (ANOVA) with orientation (upright or inverted) and TMS site (rOFA, rLO, no TMS) as independent factors revealed main effects of orientation [ $F(1,9) = 24.6, p = 0.001$ ] and of TMS site [ $F(2,18) = 12.1, p < 0.0001$ ]. Importantly there was also a significant interaction between orientation and TMS site [ $F(2,18) = 5.6, p = 0.013$ ]. Planned Bonferroni corrected tests revealed that discrimination of upright faces was significantly impaired by TMS delivered over the rOFA compared with TMS delivered over rLO ( $p = 0.008$ ) and the no TMS condition ( $p = 0.014$ ) but there was no significant difference between the rLO and no TMS condition ( $p = 0.8$ ). By contrast discrimination of inverted faces was significantly impaired by TMS delivered over rOFA ( $p = 0.044$ ) and rLO ( $p = 0.034$ ) compared with the no TMS condition.

To further demonstrate that TMS delivered over rOFA disrupted upright and inverted face discrimination while TMS delivered over rLO disrupted inverted face discrimination only we performed additional analyses to separately compare rOFA and rLO performance with the no TMS condition. As predicted, a two-by-two ANOVA examining performance in the rOFA condition with orientation (upright or inverted) and TMS site (rOFA or no TMS) as independent factors revealed a main effect of orientation [ $F(1,9) = 7.6, p = 0.022$ ] and TMS site [ $F(1,9) = 18.8, p = 0.002$ ] but no significant interaction [ $F(1,9) = 1.6, p = 0.229$ ]. By contrast a two-by-two ANOVA examining performance in the rLO condition with

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات