



# A new characterization of comonotonicity and its application in behavioral finance



Zuo Quan Xu

Department of Applied Mathematics, The Hong Kong Polytechnic University, Hong Kong

## ARTICLE INFO

### Article history:

Received 20 August 2013  
Available online 25 March 2014  
Submitted by Goong Chen

### Keywords:

Comonotonicity  
Behavioral finance  
Quantile formulation  
Atomless/non-atomic  
Pricing kernel  
Cumulative prospect theory  
Rank-dependent utility theory  
Economic equilibrium model

## ABSTRACT

It is well-known that an  $\mathbb{R}^n$ -valued random vector  $(X_1, X_2, \dots, X_n)$  is comonotonic if and only if  $(X_1, X_2, \dots, X_n)$  and  $(Q_1(U), Q_2(U), \dots, Q_n(U))$  coincide *in distribution*, for *any* random variable  $U$  uniformly distributed on the unit interval  $(0, 1)$ , where  $Q_k(\cdot)$  are the quantile functions of  $X_k$ ,  $k = 1, 2, \dots, n$ . It is natural to ask whether  $(X_1, X_2, \dots, X_n)$  and  $(Q_1(U), Q_2(U), \dots, Q_n(U))$  can coincide *almost surely* for *some* special  $U$ . In this paper, we give a positive answer to this question by construction. We then apply this result to a general behavioral investment model with a law-invariant preference measure and develop a universal framework to link the problem to its quantile formulation. We show that any optimal investment output should be anti-comonotonic with the market pricing kernel. Unlike previous studies, our approach avoids making the assumption that the pricing kernel is atomless, and consequently, we overcome one of the major difficulties encountered when one considers behavioral economic equilibrium models in which the pricing kernel is a yet-to-be-determined unknown random variable. The method is applicable to general models such as risk sharing model.

© 2014 Elsevier Inc. All rights reserved.

## 1. Introduction

The concept of comonotonicity has wide applications in actuarial science and financial risk management, see, e.g., Dhaene, Denuit, Goovaerts, Kaas, and Vyncke [5,6], Di Nunno and Øksendal [7], Deelstra, Dhaene, and Vanmaele [4]. It mainly refers to the perfect positive dependence between the components of a random vector, essentially saying that they can be represented as increasing functions of a single random variable. In this paper, we will show that these functions can be specified as their individual's quantile functions and this random variable specified as a random variable uniformly distributed on the unit interval  $(0, 1)$  that is comonotonic with the components of the random vector.

This paper consists of two parts. The first part is dedicated to studying a new characterization of comonotonic random vectors. In the second part, we apply this characterization to a general behavioral investment problem with a law-invariant preference measure and develop a universal framework to link

E-mail address: maxu@polyu.edu.hk.

the problem to its quantile formulation, which overcomes one of the major difficulties encountered when one considers a behavioral economic equilibrium model with a law-invariant preference measure where the classical dynamic programming and probabilistic approaches do not work.

We start by introducing the concept of comonotonicity for a random vector. Let us first recall the definition and the characterizations of a comonotonic random vector. For the rest part of this paper, all the random variables/vectors are in the same given probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ .

**Definition 1.** An  $\mathbb{R}^n$ -valued random vector  $(X_1, X_2, \dots, X_n)$  is comonotonic if there exists a set  $\widehat{\Omega} \times \widehat{\Omega} \subseteq \Omega \times \Omega$ , with full measure, such that

$$(X_i(\omega') - X_i(\omega))(X_j(\omega') - X_j(\omega)) \geq 0, \quad \text{for all } (\omega', \omega) \in \widehat{\Omega} \times \widehat{\Omega}, \quad i, j \in \{1, 2, \dots, n\}.$$

We denote by  $\mathbb{U}$  the set of all random variables uniformly distributed on the unit interval  $(0, 1)$  in the probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ . In order to get a new characterization of comonotonic random vector, it is necessary to assume, throughout this paper, that

**Assumption 1.** The set  $\mathbb{U}$  is not empty.

Define the quantile function  $Q_X(\cdot)$  of an  $\mathbb{R}$ -valued random variable  $X$  as the right-continuous inverse function of its cumulative distribution functions (cdf)  $F_X(\cdot)$ , that is,

$$Q_X(x) = \sup\{t \in \mathbb{R}: F_X(t) \leq x\}, \quad x \in (0, 1),$$

with convention  $\sup \emptyset = -\infty$ .

The following well-known result characterizes the comonotonic random vector (see, e.g., Dhaene, Denuit, Goovaerts, Kaas, and Vyncke [5,6]).

**Theorem 1.** An  $\mathbb{R}^n$ -valued random vector  $\mathcal{X} = (X_1, X_2, \dots, X_n)$  is comonotonic if and only if one of the following conditions holds:

1. For any vector  $\underline{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , we have

$$F_{\mathcal{X}}(\underline{x}) = \min\{F_{X_1}(x_1), F_{X_2}(x_2), \dots, F_{X_n}(x_n)\},$$

where  $F_{\mathcal{X}}(\cdot)$  and  $F_{X_k}(\cdot)$ ,  $k = 1, 2, \dots, n$ , are the cumulative distribution functions (cdf) of  $\mathcal{X}$  and  $X_k$ ,  $k = 1, 2, \dots, n$ , respectively;

2. For any  $U \in \mathbb{U}$ , we have

$$\mathcal{X} \stackrel{d}{=} (Q_1(U), Q_2(U), \dots, Q_n(U)),^1$$

where  $Q_k(\cdot)$  are the quantile functions of  $X_k$ ,  $k = 1, 2, \dots, n$ ;

3. There exist a random variable  $Y$  and non-decreasing functions  $f_k(\cdot)$ ,  $k = 1, 2, \dots, n$ , such that

$$\mathcal{X} \stackrel{d}{=} (f_1(Y), f_2(Y), \dots, f_n(Y));$$

4. Let  $Y = X_1 + X_2 + \dots + X_n$ . There exist non-decreasing functions  $f_k(\cdot)$ ,  $k = 1, 2, \dots, n$ , such that

$$\mathcal{X} = (f_1(Y), f_2(Y), \dots, f_n(Y)).^2$$

---

<sup>1</sup> We write  $X \stackrel{d}{=} Y$  if random variables/vectors  $X$  and  $Y$  are identically distributed.  
<sup>2</sup> For any two random variables/vectors  $X$  and  $Y$ , we write  $X = Y$  if  $\mathbf{P}(X = Y) = 1$ .

متن کامل مقاله

دریافت فوری ←

**ISI**Articles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات