

ScienceDirect



IFAC-PapersOnLine 48-3 (2015) 1375-1379

## The Newsvendor Problem with Fast Moving Items and a Compound Poisson Price Dependent Demand

Anna V. Kitaeva, Valentina I. Subbotina\*, Oleg A. Zmeev

National Research Tomsk State University, Tomsk, Russia, (e-mail: kit1157@yandex.ru).

**Abstract.** We consider a single period single item inventory system when the demand is a compound Poisson process with price dependent intensity and continuous batch size distribution with known mean and variance, and the mean of the demand is large enough. Equations for retail price maximizing the expected profit under optimal order quantity are obtained and approximate solution is proposed. Approximate distribution of the large order's selling time is found and results of testing procedures are given.

© 2015, IFAC (International Federation of Automatic Control) Hosting by Elsevier Ltd. All rights reserved.

Keywords: Compound Poisson, Pricing, Newsvendor, Fast Moving Items, Selling Time, Expected Profit.

## 1. INTRODUCTION AND PROBLEM STATEMENT

The newsvendor problem (NP) is one of the classical problems of inventory management; see, for example, Arrow, Harris, and Marshak (1951); Silver, Pyke, and Peterson (1998). It has been studied since the eighteenth century and widely used to analyse systems with perishable products in such different fields as, for example, health insurances, airlines, sports and fashion industries. And nowadays a lot of papers related to this problem are still being published, see reviews by Khouja (1999); Qin et al. (2011); a handbook editing by Tsan-Ming Choi (2014).

Originally the price is exogenous variable in the NP. Whitin (1955) was the first who analyzes the price-dependent demand and determines the order size and selling price simultaneously. The topic has been the theme of many papers; see reviews by Petruzzi and Dada (1999); Yano and Gilbert (2003); Chen and Simchi-Levi (2012).

Consider a supply chain consisting of a supplier, a buyer, and customers. The duration of the product's lifetime is T. At the beginning of a time period T the buyer purchases a quantity Q at a fixed price per unit d (wholesale price).

Let the demand be a Poisson process with price-dependent intensity  $\lambda(c)$ , where c > d is a selling (retail) price per unit of the product, the values of orders (batch sizes) be i.i.d. continuous random variables with finite the first and second moments equals respectively  $a_1$  and  $a_2$ . So we consider a stochastic static model.

We assume that the buyer operates with no capacity restrictions, and we do not consider the cost of leftovers utilization and lost sales. The aim is to maximize the expected profit and to find the distribution of the order's selling time. Due to complexity of the problem we solve it for fast moving items. It gives us a possibility to use the normal approximation to distribution of demand and the diffusion approximation of demand process. In such framework we manage to obtain the equation for optimal price, see (8), and the distribution of the selling time, see (13) and (14), in closed form. The latter can be used to estimate the mean and standard deviation of demand in the case of unobserved lost sales.

Note that taking into account the approximation the price optimization's task can be solved in the framework of additive-multiplicative model of demand firstly presented by Young (1978).

Denote X(t) a random customer demand at [0,t],  $p(\cdot)$  the probability density function of X(T) = X.

Expected profit for the buyer at the end of the period

$$S = -Qd + cQ\int_{Q}^{\infty} p(x)dx + c\int_{0}^{Q} xp(x)dx.$$

The buyer is interested in determining an optimal value of Q and then corresponding value of c by maximizing the expected profit. The characteristics of the duration of the order's selling are also of interest.

Obviously the first task has unique solution determined by the equation

$$\int_{Q}^{\infty} p(x)dx = \frac{d}{c},$$
(1)

and corresponding profit

$$S_0 = c \int_0^Q x p(x) dx.$$
 (2)

2405-8963 © 2015, IFAC (International Federation of Automatic Control) Hosting by Elsevier Ltd. All rights reserved. Peer review under responsibility of International Federation of Automatic Control. 10.1016/j.ifacol.2015.06.278

<sup>\*</sup>This work is performed under the state order No. 1.511.2014/K of the Ministry of Education and Science of the Russian Federation.

The result is well known; see, for example, Silver, Pyke, and Peterson (1998).

So we need to know distribution of X. In general the distribution is very complicated and it is difficult to solve the NP in closed form.

Denote probability density of the batch sizes  $f(\cdot)$ . Consider, for example, the case of exponential distribution

$$f(x) = \frac{1}{a_1} \exp\left(-\frac{x}{a_1}\right), \quad x \ge 0.$$
(3)

It is easy to see that

$$p(x) = \delta(x)e^{-\lambda(c)T} + e^{-x/a_1 - \lambda(c)T} \sqrt{\frac{\lambda(c)T}{a_1 x}} I_1\left(2\sqrt{\frac{\lambda(c)Tx}{a_1}}\right),$$

where  $I_1(\cdot)$  is the modified Bessel function of the first kind and first order,  $\delta(\cdot)$  is the Dirac delta function.

Formula (1) can be written as

$$\sqrt{\lambda(c)T} \int_{Q/a_1}^{\infty} \frac{1}{\sqrt{u}} I_1\left(2\sqrt{\lambda(c)Tu}\right) e^{-u-\lambda(c)T} du = \frac{d}{c}.$$
 (4)

Only numerical solution (4) with respect to Q is possible and the task of price optimization given Q becomes enough of an onus. So below we consider the approximate solution of the problem for fast moving items.

## 2. PRICE OPTIMIZATION FOR FAST MOVING ITEMS

Let us consider the case  $\lambda(c)T >> 1$ .

Taking into account  $I_1(z) \sim \frac{1}{\sqrt{2\pi z}} e^z$  as  $z \to \infty$ ; see Abramowitz and Stegun (1972), we receive from (4)

$$Q = a_1 \lambda(c) T + \sqrt{a_2 \lambda(c) T} \Psi\left(1 - \frac{d}{c}\right), \tag{5}$$

where 
$$\Psi(\cdot) = \Phi^{-1}(\cdot), \quad \Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} \exp\left(\frac{-t^2}{2}\right) dt$$
.

Formula (5) is valid in general, i.e. for any distribution  $f(\cdot)$ as  $\lambda T \to \infty$ , because according central limit theorem  $\frac{X - a_1\lambda(c)T}{\sqrt{a_2\lambda(c)T}}$  converges in distribution to a standard normal

random variable N(0,1).

Formula (5) is also well-known as exact result for normally distributed demand. We know that the normal distribution provides a good empirical fit to observed demand data for fast moving items.

Now find the optimal retail price for order quantity Q defined by (5).

From (2) we get as  $\lambda(c)T >> 1$ 

$$S_0 = c \left[ a_1 \lambda(c) T \Phi \left( \frac{Q - a_1 \lambda(c) T}{\sqrt{a_2 \lambda(c) T}} \right) - \sqrt{\frac{a_2 \lambda(c) T}{2\pi}} \exp \left( -\frac{(Q - a_1 \lambda(c) T)^2}{2a_2 \lambda(c) T} \right) \right]$$

For optimal value Q defined by (5) we get

$$S_0 = c \left[ a_1 \lambda(c) T \left( 1 - \frac{d}{c} \right) - \sqrt{\frac{a_2 \lambda(c) T}{2\pi}} \exp \left( -\frac{1}{2} \Psi^2 \left( 1 - \frac{d}{c} \right) \right) \right].$$

It is obvious that  $S_0 \ge 0$ , and  $S_0 \approx 0$  as  $c \approx d$ . Let the following natural condition holds:  $c\lambda(c) \approx 0$  as c >> 1, then  $S_0 \approx 0$  as c >> 1. It follows that maximum value of  $S_0(c)$  exists.

Let  $\lambda(c) = \lambda_0 F(c)$ , where  $F(\cdot) \downarrow$  is a twice differentiable function. Then

$$S_0 = a_1 \lambda_0 T \left[ F(c)(c-d) - \frac{a_2 F(c)}{2\pi a_1^2 \lambda_0 T} \exp\left(-\frac{1}{2} \Psi^2 \left(1 - \frac{d}{c}\right)\right) \right].$$
(6)

If  $\lambda(c)T >> 1$  then considering only the first summand in (6) we get a "null" approximation  $c_0$  to optimal value of retail price  $c^o$ 

$$c_0 = \arg\max_c \left( F(c)(c-d) \right).$$

The approximation is defined by equation

$$c_0 + \frac{F(c_0)}{F'(c_0)} = d.$$
 (7)

Price  $c_0$  corresponds to the case when the number of orders and, consequently, the lot size are very large.

Taking derivative 
$$\frac{dS_0}{dc}$$
 we receive the equation for  $c^o$   
 $F'(c)(c-d) + F(c) - \sqrt{\frac{a_2}{2\pi a_1^2 \lambda_0 T}} \times \left[ \left( \sqrt{F(c)} + \frac{cF'(c)}{2\sqrt{F(c)}} \right) \exp\left(-\frac{1}{2}\Psi^2\left(1 - \frac{d}{c}\right)\right) + \right]$ 

## دريافت فورى 🛶 متن كامل مقاله

- امکان دانلود نسخه تمام متن مقالات انگلیسی
   امکان دانلود نسخه ترجمه شده مقالات
   پذیرش سفارش ترجمه تخصصی
   امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
   امکان دانلود رایگان ۲ صفحه اول هر مقاله
   امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
   دانلود فوری مقاله پس از پرداخت آنلاین
   پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات
- ISIArticles مرجع مقالات تخصصی ایران