



# Relationships between posttraumatic stress disorder (PTSD), dissociation, quality of life, hopelessness, and suicidal ideation among earthquake survivors

Osman Ozdemir<sup>a</sup>, Murat Boysan<sup>b</sup>, Pinar Guzel Ozdemir<sup>a,\*</sup>, Ekrem Yilmaz<sup>a</sup>

<sup>a</sup> Yüzüncü Yıl University, Department of Psychiatry, Dursun Odabas Medical Center, Van, Turkey

<sup>b</sup> Yüzüncü Yıl University, Department of Psychology, Faculty of Arts, Van, Turkey

## ARTICLE INFO

### Article history:

Received 2 December 2014

Received in revised form

10 April 2015

Accepted 3 May 2015

Available online 10 June 2015

### Keywords:

Posttraumatic syndrome

Dissociative disorders

Depression

Suicidality

Well-being

Disability

## ABSTRACT

Researches have demonstrated that Posttraumatic stress disorder (PTSD) is one of the most common stress reactions in the face of disasters and significantly associated with a broad range of trauma-induced sequelae including anxiety, depression, suicidality as well as functional impairments. To date, though many aspects of risk factors with respect to the development and maintenance of PTSD have been addressed, mediating role of dissociation has received relatively less attention. In the present study, we examined relations of PTSD with quality of life, hopelessness, suicidal ideation, and mediational effect of pathological dissociation in these connections. 583 subjects most of whom experienced a severe earthquake participated in the study after two years of the disaster. We found that being female, being single, earthquake exposure, and having greater suicidal ideation were significant predictors of PTSD symptom severity. Role-Physical, Bodily-Pain, General Health and Role-Emotional subscales of the SF-36 were inversely associated with PTSD symptom severity. Pathological dissociation significantly mediated the substantial associations between predictors and PTSD symptom clusters. Chronic dissociation appears to put trauma exposed individuals in jeopardy of prolonged posttraumatic reactions by mediating the negative influences of risk factors in the face of experienced earthquake.

© 2015 Elsevier Ireland Ltd. All rights reserved.

## 1. Introduction

Epidemiological studies have shown that exposure to disasters i.e. earthquakes, floods, or hurricanes etc. are not rare among people and result in a broad range of mental and physical health consequences (Norris et al., 2002). Posttraumatic stress disorder (PTSD) is one of the most common stress reactions and presumably the most debilitating condition in case of trauma exposure, specifically disasters (Galea et al., 2005). Most individuals exposed to aversive life events experience a natural recovery process in the aftermath of the trauma and emotional regulation comes back to a level prior to the incident. Nevertheless, a substantial minority fail to extinguish eliciting various stress responses that perpetuate trauma-induced acute reactions in chronic suffering from severe symptoms, particularly PTSD (Breslau, 2002; Kessler et al., 1995).

Although defining the construct of dissociative phenomena is a matter of various debates because of its multifaceted characteristics, trauma-induced dissociation can be defined as a disruption in the

usually integrated functions of mental processes including sense perceptions, thoughts, feelings, experiences, and memories (DePrince and Freyd, 2007). Significant linkages between the trauma and dissociation have also been well-established. Research has shown that dissociative symptoms have an important role in the development and maintenance of posttraumatic stress disorder (PTSD) (Galatzer-Levy et al., 2011; Stein et al., 2013; Wolf et al., 2012). Nonetheless, co-occurring dissociation is not peculiar to PTSD rather a broad range of dissociative symptomatology could be commonly observed among patients with other psychiatric disorders and even in general population (Giesbrecht et al., 2004; Maaranen et al., 2005; Mulder et al., 1998). Patients with dissociative symptoms have significantly higher rates of comorbid psychiatric conditions including depression, borderline personality disorder, and self-mutilation and suicide attempts (Ebrinc et al., 2008; Maaranen et al., 2005; Zoroglu et al., 2003). Also, it has been consistently demonstrated that dissociative experiences were associated with impaired quality of life such as physical and social functioning (Mitchell et al., 2012).

Bremner (1999) proposed a dualistic model of posttraumatic reactions is that psychological manifestations in response to traumatic events can be best understood by grouping posttraumatic syndrome into two lines: one predominantly dissociative and the other characterized by hyperarousal states. Experiencing

\* Correspondence to: Yüzüncü Yıl University, Faculty of Medicine, Department of Psychiatry, Van 65200, TURKEY. Tel.: +90 4322150474, fax: +90 4322168352.

E-mail address: [pguzelozdemir@yahoo.com](mailto:pguzelozdemir@yahoo.com) (P. Guzel Ozdemir).

dissociative symptoms during the adverse events strongly predicts latter onset of persistent PTSD (Friedman, 2013; Ozer et al., 2003) and dissociative symptoms such as numbing are a necessary condition for acute and posttraumatic stress disorder diagnosis in DSM-5 (American Psychiatric Association, 2013). Due to the stress–diathesis model of dissociation, dissociative mechanisms are implicated in either buffering stress-related extreme responses or disrupting affect regulation (Butler et al., 1996). Briere et al. (2005) submitted a distinction between peritraumatic and chronic pathological dissociation that, in comparison to the former, the latter was likely to represent a more pathological subtype of dissociation with worse psychological outcomes. Recent advances in PTSD research have provided supportive evidence for that dissociative symptomatology central to PTSD; whereas a sizable minority of individuals with persistent PTSD at about 30% seem to have a tendency to elicit more severe dissociative symptoms relative to the other subset of PTSD patients experiencing primarily hyperarousal states with relatively low dissociative inclination in response to reminders of trauma memory (Lanius et al., 2012). Moreover, the premise that dissociative phenomena convey both buffering and maladaptive effects among trauma exposed person was plainly demonstrated in a representative sample by Boysan (2014) that normal dissociation characterized by lower levels of dissociative experiences was negatively associated with measures of detrimental affective states; on the other extreme, greater levels of dissociative symptomatology was a significant predictor of elevations in affective symptoms. Over and above, pathological dissociation was not only a significant antecedent for persistent PTSD but also a mediator of significant associations of PTSD symptoms with other predictors as a risk factor (Kadak et al., 2013; Özdemir et al., *In press*).

In the epidemiological studies of psychopathology, PTSD has exhibited an excessively discrete pattern of comorbidity relative to other psychiatric disorders and consistently detected to be co-occurring with other nosological entities to an extent to a higher of 92% (Keane et al., 2007). Significant linkages between PTSD and suicidal behaviors have long been recognized and an increase in research interest focusing on these relation has been carried on. Comorbid psychiatric conditions have been found to be conferring PTSD affected people at higher risk of suicidality across different populations, such as war veterans, survivors of disasters, victims of childhood and/or adult trauma (Krysinska and Lester, 2010; Panagioti et al., 2009; 2012). Some authors have postulated that traumatization result in disruption of basic psychobiological constructs that constitute personality (Hulet et al., 2011; Mauritz et al., 2013). These formulations suggest that traumatized individuals could exhibit behavioral problems, such as impulsivity, aggression, substance abuse, and self-injurious acts, as well as internalization problems including symptoms of depression and anxiety (Stein et al., 2013). In addition, commonly found was specifically major depression which seems to have a central role in a tendency to suicidal behaviors. Leiner et al. (2008) evidenced for mediational influence of comorbid depressive disorder in accounting the substantial relations between suicidal ideation and PTSD. A comprehensive meta-analysis put forth a series of considerable suggestions that significant correlates of PTSD were suicidal ideation and suicidal behaviors rather than completed suicides, and these relations were sustained irrespective of subtypes of traumatic experiences and of samples recruited from either psychiatric or community populations (Panagioti et al., 2012).

Despite suicide has long being a matter of intense interest and the extensive clinical and epidemiological literature, there are still huge gaps in our understanding of factors associated with suicidality and suicidal thought (Silverman, 2011). To the best of our knowledge, suicidality may be a relatively stable vulnerability for some individuals and unlike suicidal behaviors, suicidal ideation

by its own may be a specific cognitive susceptibility factor for development of psychopathology rather than co-occurring with psychopathology, such as depression or PTSD. Poor problem solving capabilities, greater impulsivity (Windfuhr and Kapur, 2011), and deficiencies in interpersonal skills, conceptualized as burdensomeness and thwarted relatedness (Ribeiro and Joiner, 2011), have been suggested to be the hallmark features of suicidal individuals. Suicidal behavior is one of pathological manifestations which represents incapability in dealing with stressful situations coupled with emotional regulation deficits. General coping deficiencies in conjunction with emotional dysregulation may lead to both a failure in extinguishing acute stress responses and suicidal behaviors along with the psychopathology in the face of traumatic events. Turning to higher order cognitive information processing in suicide, the perception of defeat and entrapment are speculated to be significant correlates of suicidal behaviors (Johnson et al., 2008) which seem reminiscent of the conceptualization of mental defeat in the cognitive model of PTSD proposed by Ehlers and Clark (2000). Overlapping cognitive mechanisms in suicide and PTSD still remain elusive and need further investigations.

To date, several models have been proposed to explain the relationship between dissociation, and self-destructive behaviors that are important in suicide attempts. First, self mutilation might be related to regulate overwhelming discomfort of dissociative experiences, sensations, emotions or feelings resulting from immature and maladaptive coping strategies (Chang et al., 2010). Second, dissociative mechanisms may be upholding the pain threshold and pain tolerance during the self-injury behaviors (Orbach, 1994). Third, dissociative symptoms often accompany to psychiatric disorders, particularly major depression, and borderline personality disorder in which higher rates of suicidal behaviors are observed (Chang et al., 2010; Grabe et al., 1999; Klonsky and Moyer, 2008). Although, suicidal thoughts and proneness to suicidal behaviors generally suggested to be a function of mood regulation problems and psychopathology, to the best of our knowledge, these relations are likely to be reciprocally determined that suicidality may also be an important antecedent of mood symptoms and self-regulation problems. In a preliminary study, Selvi et al. (2010) demonstrated that, by using multi-sample path modeling, suicidal ideation was not predicted by depressive symptomatology but rather depressive symptoms were a product of suicidal ideation. In this vein, suicidality may be the antecedent of PTSD as a function of mediational effect of pathological dissociation.

In a recent study conducted amongst a representative community sample, Sareen et al. (2007) plainly provided convincing evidence for that PTSD remains a significant predictor of physical health problems after controlling for socio-demographic features and other mental disorders. PTSD was also predictive of suicide attempts, poor quality of life, and short- and long-term disability after adjusting for socio-demographic factors, mental disorders, and severity of physical disorders. Previous studies have consistently reported poor health status and functional impairment among disaster survivors. Lai et al. (2004) examined the differences in a set of demographic and clinical features between full, partial and non-PTSD groups after 10 months of a severe earthquake in Taiwan and found that individuals with either full or partial PTSD reported significantly greater levels of suicidality, general psychopathology, disability; and impaired well-being relative to individuals with non-PTSD. Injured victims of earthquake reported higher levels of PTSD symptoms compared to individuals with no injury due to the disaster (Kuo et al., 2007). Six months after the devastating floods and mudslides in Mexico, acute PTSD reactions were found to mediate the relations between disaster exposure and physical health problems in a sample of adults. Regardless of whether physical health was medically

متن کامل مقاله

دریافت فوری ←

**ISIArticles**

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلید کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات