

Water management in the framework of environmental management systems in Bulgarian seaports

Le Xuan Quynh^{a,*}, Luc Hens^a, Stoyan Stoyanov^{b,1}

^a Department of Human Ecology, Vrije Universiteit Brussel, Laarbeeklaan 103, B-1090 Brussels, Belgium

^b Ecology Center, University of Chemical Technology & Metallurgy, 8 Kl. Ohridski blvd, 1756 Sofia, Bulgaria

ARTICLE INFO

Article history:

Received 12 January 2009

Received in revised form 28 April 2010

Accepted 15 May 2010

Available online 21 May 2010

Keywords:

EMS

Seaport

Environmental policy

Significant environmental aspect

Bourgas

Varna

ABSTRACT

Seaports or harbours are major hub of economic activities, connecting sea routes with the hinterland via rail, road and inland waterway. At the same time, they are hubs of environmental issues, resulting from its diversified operations involving vessels, machines, vehicles and industries. The Black Sea Commission in 2007 pointed out that water pollution at the ports around Black Sea, including the ports of Bourgas and Varna, poses a great threat to the overall water quality of the Black Sea and the quality of the Mediterranean Sea. The two ports, and their facilities, are the black spots of pollution that need immediate intervention to safeguard the Black Sea. The first steps of an environmental management system were implemented at both ports. Environmental protection objectives are identified and initial actions are being implemented. However, assessment of the environmental performance of both ports based on concrete indicators was not done. Further efforts are needed to advance the environmental management system to contribute more to the protection of the Mediterranean and Black Sea region.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

Seaports are major hubs of economic activities. It is an increasing trend that they handle not only ships but they are crossroads of transport networks where sea routes meet rail, road and inland navigation and vice versa (ESPO, 2004; ESPO-ITMMA, 2007). Seaports are the main gateway for Europe with constant movement of ships, trains, trucks and barges, bringing containers, pallets, bags, and bulk to and from the European hinterland. On average, European harbours accommodate 2 million vessels per year and handle 3.5 billion tonnes of goods and 350 million passengers (ESPO, 2004). There are over 1000 harbours along the European coastlines (EC, 2007).

Seaports around the world face increasing pressure to develop larger and more efficient to accommodate soaring waterborne traffic. In Europe, ports are even under higher pressure to follow stringent environmental standards while maintaining their competitiveness in a very demanding market. Although maritime transports are reported to produce 13 times less pollution than road transport in terms of carbon and particulate emissions and 19 times less in terms of hydrocarbons, environmental issues at the port areas are still of great concern (ESPO, 2004). Maritime

transport has adverse environmental impacts, such as water pollution through the discharges of oil and sewage water from ports and vessels, air pollution from exhaust gases emitted from the vessel's engine, water pollution by toxins (such as Tributyltin or TBT and its derivatives, heavy metals), and the translocation of harmful organisms and pathogens via ballast water and sediments inside ballast water tanks (Hallegraeff and Bolch, 1991; Gollasch, 2006; Gollasch and Leppäkoski, 2007; Ruiz et al., 2000; David et al., 2007a,b; EU Commission, 2007; UNCTAD, 2007).

The Mediterranean and the Black Sea regions host the highest density of ports in Europe. Amongst 180 main ports that together handling at least 80% of the country's cargo traffic in the whole of Europe-25 in 2004, 63 are located around the Mediterranean Sea (EC, 2007). (Note: Main ports are those handle over 1 million tonnes of goods per year.) This figure excludes ports in Turkey, Georgia, Ukraine and Russia around the Black Sea and other Mediterranean countries that not belong to the European Union.

Bulgarian seaports face many environmental issues such as air pollution, water pollution, dredging, wastes, oily waste, etc. They are classified as "black points" of pollution in the Black Sea region that require special attention and actions (BSC, 2002, 2007). Therefore, there is an urgent need to put an Environmental Management System (EMS) in place to deal with those problems.

An EMS is a continuous cycle of planning, implementation, review and improvement of processes and actions that an organisation undertakes to meet its environmental obligations. It is a part of the overall management system and contains all components

* Corresponding author. Tel.: +32 2 477 4281; fax: +32 2 477 4964.

E-mail addresses: Le.XuanQuynh@vub.ac.be (L.X. Quynh), stoyan1@uctm.edu (S. Stoyanov).

¹ Tel.: +359 2 868 1021.




Fig. 1. Deming cycle and the elements of EMS.

reducing environmental liability; managing effectively resources in achieving environmental obligations; and improving general business/production performance (Fig. 1).

For the majority of the organisations, many elements of an EMS may already be in place as they are common to management processes, such as quality, health and safety, finance, or human resources. Integrating environmental management with other key processes can improve financial, quality and environmental performance. An EMS can be an investment in long-term development of the business: businesses can build a better reputation that helps to keep existing customers, clients and investors and attract new ones.

2. Materials and methods

2.1. Literature review

Data on ports' operation is collected from the ports and other published data sources using a questionnaire as a guidance on the types of information needed to be considered during the inventory as well as the format of the inventory. The questionnaire consisted of eight elements: 1. The port and its environment; 2. The port's environmental policy; 3. The use of environmental management instruments in the port; 4. Port management system; 5. Environmental legislation; 6. Land use and land use changes; 7. Significant environmental aspects; and 8. Environmental performance. For each elements from 1 to 6, data was collected on a set of indicators (Table 1).

common for a management system (Sheldon and Yoxon, 2002; Whitelaw, 2004).

The EMS has different targets: improving environmental performance; complying with environmental laws and regulations;

Table 1
Selected indicators used in the inventory.

| Aspect | Indicators |
|---|---|
| <i>Description about the port</i> | |
| | Geographical location Port's detail: name, number of berth, total berth length (m), capacity (max receivable vessel – tonnage) Production (2000–2005): cargo throughput (million tonnes), container throughput (TEUs), passenger (people) Port's facilities and human resources Land use (ha): area of warehouse, storage Coastal area around the port: type of coastline, ecosystems |
| <i>Environmental conditions in and around the ports</i> | |
| Water quality | Physio-chemical parameters: Temperature (°C), turbidity (NTU), TDS and TSS (mg/l) Organic and inorganic pollution: DO, BOD, COD, nitrite, nitrate, ammoniac, phosphate, and coliforms (MPN/100 ml) Metals in water (mg/l): Chromium (Cr), Cadmium (Cd), Copper (Cu), Lead (Pb), Manganese (Mn), Zinc (Zn), Iron (Fe), Arsenic (As), Mercury (Hg), Oil (mg/l) Persistent Organic Compounds (µg/l): lindane, aldrin, dieldrin, endrin, DDT and metabolites (DDD and DDE), organotins (TBT and derivatives) |
| Soil and sediment quality | Metals in sediments (water bottom) (mg/kg dry): Cadmium (Cd), Copper (Cu), Lead (Pb), Zinc (Zn), Arsenic (As), Mercury (Hg) Oil (mg/g dry) and cyanide (mg/kg dry) Persistent Organic Compounds (µg/kg dry weight): lindane, aldrin, dieldrin, endrin, DDT and metabolites (DDD and DDE), organotins (TBT and derivatives) |
| Air quality | Micro-climate conditions: Temperature (°C), humidity (%), Wind velocity (m/s), CO, NO ₂ , SO ₂ , Pb, VOC, dusts (mg/m ³) Noise (dBA) |
| Biological resources | Degradation of ecosystems Presence of pollutant in biological indicative species (clam, mussels, fish and resident bird) (mg/kg dry tissue): metals, persistent organic compounds |
| <i>Emissions from the port</i> | |
| Into the air | CO, CO ₂ , NO _x , SO ₂ , VOC, dust, PM (tonnes/year) |
| Into the water | Amount of household wastewater (m ³ /year) Amount of oily waste from ship (m ³ /year): bilge, oil sludge |
| Solid waste | From ships and port's operations: type; amount (tonnes/year) |
| <i>Environmental legislation related to port</i> | |
| | International conventions National laws, regulations, action plans Sectoral regulations and plans Port's policy, strategy |
| <i>Environmental management at the port</i> | |
| | Environmental Management System Risk Assessment Environmental Impact Assessment |

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات