


Author's Accepted Manuscript

Assessing and Minimizing Contamination in Time of Flight Based Validation Data

Kristin P. Lennox, Paul Rosenfield, Brenton Blair, Alan Kaplan, Jaime Ruz, Andrew Glenn, Ronald Wurtz



www.elsevier.com/locate/nima

PII: S0168-9002(17)30473-4
DOI: <http://dx.doi.org/10.1016/j.nima.2017.04.022>
Reference: NIMA59810

To appear in: *Nuclear Inst. and Methods in Physics Research, A*

Received date: 8 March 2017
Revised date: 10 April 2017
Accepted date: 15 April 2017

Cite this article as: Kristin P. Lennox, Paul Rosenfield, Brenton Blair, Alar Kaplan, Jaime Ruz, Andrew Glenn and Ronald Wurtz, Assessing and Minimizing Contamination in Time of Flight Based Validation Data, *Nuclear Inst. and Methods in Physics Research, A* <http://dx.doi.org/10.1016/j.nima.2017.04.022>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Assessing and Minimizing Contamination in Time of Flight Based Validation Data

Kristin P. Lennox^{a,1}, Paul Rosenfield^a, Brenton Blair^a, Alan Kaplan^a, Jaime Ruz^b, Andrew Glenn^b, Ronald Wurtz^b

^a*Computational Engineering Division, Lawrence Livermore National Laboratory, Livermore, California*

^b*Physics Division, Lawrence Livermore National Laboratory, Livermore, California*

Abstract

Time of flight experiments are the gold standard method for generating labeled training and testing data for the neutron/gamma pulse shape discrimination problem. As the popularity of supervised classification methods increases in this field, there will also be increasing reliance on time of flight data for algorithm development and evaluation. However, time of flight experiments are subject to various sources of contamination that lead to neutron and gamma pulses being mislabeled. Such labeling errors have a detrimental effect on classification algorithm training and testing, and should therefore be minimized. This paper presents a method for identifying minimally contaminated data sets from time of flight experiments and estimating the residual contamination rate. This method leverages statistical models describing neutron and gamma travel time distributions and is easily implemented using existing statistical software. The method produces a set of optimal intervals that balance the trade-off between interval size and nuisance particle contamination, and its use is demonstrated on a time of flight data set for Cf-252. The particular properties of the optimal intervals for the demonstration data are explored in detail.

Keywords: time of flight, pulse shape discrimination, Poisson regression,

*Corresponding author

Email address: `blair28@llnl.gov` (Brenton Blair)

¹This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات