

Accepted Manuscript

Lignocentric analysis of a carbohydrate-producing lignocellulosic biorefinery process

Robert H. Narron, Qiang Han, Sunkyu Park, Hou-min Chang, Hasan Jameel

PII: S0960-8524(17)30882-9
DOI: <http://dx.doi.org/10.1016/j.biortech.2017.05.207>
Reference: BITE 18233

To appear in: *Bioresource Technology*

Received Date: 3 April 2017
Revised Date: 30 May 2017
Accepted Date: 31 May 2017

Please cite this article as: Narron, R.H., Han, Q., Park, S., Chang, H-m., Jameel, H., Lignocentric analysis of a carbohydrate-producing lignocellulosic biorefinery process, *Bioresource Technology* (2017), doi: <http://dx.doi.org/10.1016/j.biortech.2017.05.207>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Lignocentric analysis of a carbohydrate-producing lignocellulosic biorefinery process

Robert H. Narron, Qiang Han, Sunkyu Park, Hou-min Chang, Hasan Jameel*

Department of Forest Biomaterials, North Carolina State University,

2820 Faucette Dr, Raleigh, NC, 27607

Abstract

A biologically-based lignocellulosic biorefinery process for obtaining carbohydrates from raw biomass was investigated across six diverse biomasses (three hardwoods & three nonwoods) for the purpose of decoding lignin's influence on sugar production.

Acknowledging that lignin could positively alter the economics of an entire process if valorized appropriately, we sought to correlate the chemical properties of lignin within the process to the traditional metrics associated with carbohydrate production- cellulolytic digestibility and total sugar recovery. Based on raw carbohydrate, enzymatic recovery ranged from 40-64% w/w and total recovery ranged from 70-87% w/w. Using nitrobenzene oxidation to quantify non-condensed lignin structures, it was found that raw hardwoods bearing increasing non-condensed S/V ratios (2.5-5.1) render increasing total carbohydrate recovery from hardwood biomasses. This finding indicates that the chemical structure of hardwood lignin influences the investigated biorefinery process' ability to generate carbohydrates from a given raw hardwood feedstock.

Key Words: Autohydrolysis, autohydrolyzate, sugar recovery, enzymatic digestibility, non-condensed lignin

1. Introduction

* Corresponding author, jameel@ncsu.edu

متن کامل مقاله

دریافت فوری ←

ISIArticles

مرجع مقالات تخصصی ایران

- ✓ امکان دانلود نسخه تمام متن مقالات انگلیسی
- ✓ امکان دانلود نسخه ترجمه شده مقالات
- ✓ پذیرش سفارش ترجمه تخصصی
- ✓ امکان جستجو در آرشیو جامعی از صدها موضوع و هزاران مقاله
- ✓ امکان دانلود رایگان ۲ صفحه اول هر مقاله
- ✓ امکان پرداخت اینترنتی با کلیه کارت های عضو شتاب
- ✓ دانلود فوری مقاله پس از پرداخت آنلاین
- ✓ پشتیبانی کامل خرید با بهره مندی از سیستم هوشمند رهگیری سفارشات